



Postoperative care after oral surgery

MKA

Information for patients who have had surgery in the outpatient clinic of the department of oral and maxillofacial surgery.

Eating, drinking, smoking and alcohol

When the local anesthetic has worn off, you can eat and drink as long as it is not very hot or cold. Alcohol and smoking have adverse effects on the wound healing and may increase post-operative pain. Do reduce the use of these to a minimum.

Postoperative pain, swelling and fever

Pain and swelling are normal post-operative characteristics and may last for about five days. Paracetamol is usually a sufficient painkiller and is easily available at drugstores and pharmacies. If necessary, the oral surgeon will prescribe another painkiller. The local anesthesia has worn off after two to four hours. You should better start to use the painkillers before the local anesthesia has worn off. A swollen cheek, a bluish coloration of the skin and a decreased mouth opening are normal features for a couple of days. To reduce the swelling, the face can be cooled the first day with cold packages, for example ice-cubes in a small plastic bag wrapped in a washing-glove. Fever, up to 38,5° celcius , is normal. Please contact the department if the body temperature rises over 38,5° Celsius or the fever continues for more than five days.

Bleeding

If bleeding continues after treatment, you can try to stop this by applying a compress, for example a folded gauze, on the wound. By applying gentle pressure through biting or pushing your finger for half an our, the bleeding will stop in most cases. If not, repeat the procedure. If this again does not work, contact the department.

Oral hygiene

Blood clotting in the wound is essential for problem-free wound healing. Do not rinse the mouth during the first 24 hours. The mouth can be cleaned as usual after 24 hours. Do not brush teeth at painful spots, but rinse the area with lukewarm water (1/2 spoon salt/glass).

Sutures

Usually resorbable sutures are used. If not, an appointment for a control-visit will be made. Sometimes a medicinal gauze is inserted in the wound. In that case a control-visit for removal is necessary.

Apex resection

If a temporary restoration is used in the tooth crown, replacement by a definitive restoration is required. Contact your dentist within two to four weeks after surgery for this treatment.

Sinus Perforation

Sometimes an opening between the mouth and the sinus has to be closed with non-resorbable sutures. These wounds are prone to problems in healing. Therefore it is important to avoid differences in pressure between the mouth and the nasal cavity. So, keep your mouth open during sneezing and do not blow your nose. The first 48 hours some blood may come out of the nose. This is normal and will stop by itself.

Locatie Enkhuizen

Molenweg 9b
1601 SR Enkhuizen
T. 0228 312 345

Locatie Hoorn

Maelsonstraat 3
1624 NP Hoorn
T. 0229 257 257

Locatie Purmerend

Waterlandlaan 250
1441 RN Purmerend
T. 0299 457 457

Locatie Volendam

Heideweg 1b
1132 DA Volendam
T. 0229 257 500

Postbus 600, 1620 AR Hoorn

www.dijklander.nl
info@dijklander.nl



FLD-01150-NL 10-12-2020